LACTATED RINGER'S

Injection, USP

EACH 100 mL CONTAINS SODIUM CHLORIDE 600 mg; SODIUM LACTATE, ANHYD. 310 mg; POTASSIUM CHLORIDE 30 mg; CALCIUM CHLORIDE, DIHYDRATE 20 mg IN WATER FOR INJECTION. MAY CONTAIN HCI OR NaOH FOR pH ADJUSTMENT. ELECTROLYTES PER 1000 mL (NOT INCLUDING pH ADJUSTMENT): SODIUM 130 mEg; POTASSIUM 4 mEq: CALCIUM 3 mEq: CHLORIDE 109 mEq: LACTATE 28 mEq. 273 mOsmol/LITER (CALC), pH 6.6 (6.0 to 7.5), CAUTION: DO NOT ADMINISTER CALCIUM CONTAINING SOLUTIONS CONCURRENTLY WITH STORED BLOOD, NOT FOR USE IN THE TREATMENT OF LACTIC ACIDOSIS. ADDITIVES MAY BE INCOMPATIBLE. CONSULT WITH PHARMACIST, IF AVAILABLE, WHEN INTRODUCING ADDITIVES. USE ASEPTIC TECHNIQUE, MIX THOROUGHLY AND DO NOT STORE. SINGLE-DOSE CONTAINER. FOR I.V. OR SUBCUTANEOUS USE. USUAL DOSAGE: SEE INSERT. STERILE, NONPYROGENIC, USE ONLY IF SOLUTION IS CLEAR AND CONTAINER IS UNDAMAGED. MUST NOT BE USED IN SERIES CONNECTIONS.

Rx ONLY



icumedical

IM-4294

ICU Medical, Inc., Lake Forest, Illinois, 60045 USA