

1000 mL

NDC 0990-4179-05

Aminosyn® -PF 10%

— 1

Sulfite-Free

AN AMINO ACID INJECTION — PEDIATRIC FORMULA

— 2

**Pharmacy Bulk Package —
Not for Direct Infusion.**

EACH 100 mL CONTAINS: TOTAL AMINO ACIDS APPROX. 10 g.
ESSENTIAL AMINO ACIDS/100 mL: ISOLEUCINE 760 mg; LEUCINE 1200 mg; LYSINE (AS ACETATE SALT) 677 mg; METHIONINE 180 mg; PHENYLALANINE 427 mg; THREONINE 512 mg; TRYPTOPHAN 180 mg; VALINE 673 mg. **NONESSENTIAL AMINO ACIDS/100 mL:** TYROSINE 44 mg; ALANINE 698 mg; ARGININE 1227 mg; GLYCINE 385 mg; PROLINE 812 mg; HISTIDINE 312 mg; SERINE 495 mg; GLUTAMIC ACID 820 mg; ASPARTIC ACID 527 mg; TAURINE 70 mg. **ELECTROLYTES (mEq/LITER):** ACETATE 46 mEq. 788 mOsmol/L SPECIFIC GRAVITY = 1.03 pH 5.5 (5.0 TO 6.5)
 FOR USE ONLY IN A PHARMACY ADMIXTURE PROGRAM. FOR I.V. USE. USUAL DOSAGE: SEE INSERT. STERILE, NONPYROGENIC. CONTAINS NO BACTERIOSTAT. SEE INSERT FOR COMPLETE PRODUCT INFORMATION FOR USE OF THE PHARMACY BULK PACKAGE.

— 3

— 4

— 5

DATE ENTERED: _____

TIME OF ENTRY: _____

— 6

CAUTION: USE ONLY IN LAMINAR FLOW HOOD. **ONCE THE OUTLET SITE HAS BEEN ENTERED, THE WITHDRAWAL OF CONTAINER CONTENTS SHOULD BE PROMPTLY COMPLETED IN ONE CONTINUOUS OPERATION.** DISCARD CONTAINER NOT LATER THAN 4 HOURS AFTER INITIAL CLOSURE PUNCTURE. ADDITIVES MAY BE INCOMPATIBLE WITH FLUID WITHDRAWN FROM THIS CONTAINER. CONSULT WITH PHARMACIST, IF AVAILABLE. WHEN COMPOUNDING ADMIXTURES, USE ASEPTIC TECHNIQUE. MIX THOROUGHLY. DO NOT STORE SOLUTIONS CONTAINING ADDITIVES. BECAUSE OF THE POTENTIAL FOR LIFE-THREATENING EVENTS, CAUTION SHOULD BE TAKEN TO ENSURE THAT PRECIPITATES HAVE NOT FORMED IN ANY PARENTERAL NUTRIENT MIXTURE. STORE AT 20 TO 25°C (68 TO 77°F). [SEE USP CONTROLLED ROOM TEMPERATURE.] AVOID EXCESSIVE HEAT. PROTECT FROM FREEZING. AVOID EXPOSURE TO LIGHT. USE ONLY IF SOLUTION IS CLEAR AND CONTAINER IS UNDAMAGED. MUST NOT BE USED IN SERIES CONNECTIONS.

— 7

— 8

Rx ONLY



CONTAINS DEHP

IM-4396

icumedical

— 9

ICU Medical, Inc., Lake Forest, Illinois, 60045, USA



(01)00309904179059