Lactated Ringer's Injection, USP

01)0030990795303

FACH CONTAINS SODIUM CHLORIDE 600 mg; SODIUM LACTATE. ANHYD. 310 mg; POTASSIUM CHLORIDE 30 mg; CALCIUM CHLORIDE, DIHYDRATE WATER 20 FOR INJECTION. CONTAIN HCL 0R MAY Na_OH ADJUSTMENT. **ELECTROLYTES** PER 1000 mL (NOT INCLUDING pH ADJUSTMENT): SODIUM 130 POTASSIUM 4 mEg; CALCIUM 3 mEg; CHLORIDE 109 mEq: LACTATE 28 mEq. 273 mOsmol/LITER (CALC). pH 6.6 (6.0 to 7.5). CAUTION: DO NOT ADMINISTER CONTAINING SOLUTIONS CONCURRENTLY WITH STORED BLOOD. NOT FOR USE IN THE TREATMENT LACTIC ACIDOSIS. ADDITIVES MAY INCOMPATIBLE. CONSULT PHARMACIST. IF AVAILABLE. INTRODUCING ADDITIVES, USE ASEPTIC TECHNIQUE, MIX THOROUGHLY AND DO NOT STORE. SINGLE-DOSE CONTAINER. FOR I.V. OR SUBCUTANEOUS USE. USUAL DOSAGE: SEE INSERT. NONPYROGENIC. USE ONLY IF SOLUTION AND CONTAINER UNDAMAGED. MUST NOT BE USED IN SERIES CONNECTIONS.

<u>50</u>

100

150

200



Rx only



icumedical

IM-4359